

REGULARITY OF COBOUNDARIES FOR NON UNIFORMLY EXPANDING MARKOV MAPS

SÉBASTIEN GOUËZEL

ABSTRACT. We prove that solutions u of the equation $f = u - u \circ T$ are automatically Hölder continuous when f is Hölder continuous, and T is non uniformly expanding and Markov. This result applies in particular to Young towers and to intermittent maps.

1. RESULTS

Let (X, m) be a probability space and $T : X \rightarrow X$ an ergodic measure preserving transformation. Let also G be a locally compact abelian group, endowed with an invariant metric that we denote by $|x - y|$. It is often important to know whether a function $f : X \rightarrow G$ is a *measurable coboundary*, i.e., there exists a measurable function $u : X \rightarrow G$ such that

$$(1) \quad f = u - u \circ T$$

almost everywhere. For $G = \mathbb{R}$, this condition is indeed often the only obstruction to have a non-degenerate central limit theorem for the Birkhoff sums of f (see e.g. [Leo60], [GH88], [Liv96]). For $G = S^1$, it is relevant to prove local limit theorems (see [AD01] and [ADSZ04] when f is locally constant, in the Markov and non-Markov case).

When T is uniformly hyperbolic and f is Hölder continuous, the Livšic regularity theorem ([Liv72]) states that u must have a Hölder continuous version, for which (1) holds everywhere. In particular, if there exists a point x such that $T^n(x) = x$ and $\sum_{k=0}^{n-1} f(T^k x) \neq 0$, then f is not a measurable coboundary. Hence, it is possible to prove in practice that a function is not a coboundary (see also [PY99] and [NS03]).

In this note, we extend the aforementioned result of Livšic to non-uniformly expanding Markov dynamical systems, without any additional assumption on the functions f or u . The result will first be given in the abstract setting of *Gibbs-Markov maps* (see [Aar97]). Applications to Young towers, intermittent maps in dimension 1 and positive recurrent Markov shifts will also be described.

The proof is quite flexible since it is completely elementary and does not use spectral theory. Hence, the same kind of arguments may be used in other settings.

1.1. Results for Gibbs-Markov maps. In this paragraph, we will work in the setting of Gibbs-Markov maps, defined in [Aar97, Section 4.7].

Let us recall briefly the definitions. Let $(X, d, \mathcal{B}, m)$ be a bounded metric space endowed with its Borel σ -algebra and a probability measure. A non singular map

Date: January 25, 2005.

2000 Mathematics Subject Classification. 37A20, 37D25.

$T : X \rightarrow X$ is *Gibbs-Markov* if there exists a partition α of X (modulo 0) by sets of positive measure, such that

- (1) For all $a \in \alpha$, $T(a)$ is a union (modulo 0) of elements of α and $T : a \rightarrow T(a)$ is invertible.
- (2) There exists a finite subset $\{a_1, \dots, a_n\}$ of α with the following property: for any $a \in \alpha$, there exist $i, j \in \{1, \dots, n\}$ such that $a \subset T(a_i)$ and $a_j \subset T(a)$ (modulo 0).
- (3) Expansion: there exists $\lambda > 1$ such that $\forall a \in \alpha$, for almost all $x, y \in a$, $d(Tx, Ty) \geq \lambda d(x, y)$.
- (4) Distortion: for $a \in \alpha$, let g be the inverse of the jacobian of T on a , i.e., $g(x) = \frac{dm|_a}{d(m \circ T|_a)}(x)$ for $x \in a$. Then there exists C such that, for all $a \in \alpha$, for almost all $x, y \in a$, $\left|1 - \frac{g(x)}{g(y)}\right| \leq Cd(Tx, Ty)$.

Property (2), also known as the BIP (big images and preimages) property, is apparently stronger than the usual *big image property* $\inf_{a \in \alpha} m(Ta) > 0$. However, when (4) is satisfied and T is probability preserving, these two properties are equivalent by [Sar03].

Usually, Gibbs-Markov maps are endowed with a distance given by $d(x, y) = \tau^{s(x, y)}$ where $\tau \in (0, 1)$ and $s(x, y)$ is the separation time of x and y . We have chosen here to use a general distance since it will be more convenient in the applications: our main result will say that a function is Lipschitz continuous with respect to d , which means that having more freedom to choose the distance will give more precise results. In particular, when the Gibbs-Markov map is obtained by coding another dynamical systems, it is natural to use the distance induced by the original distance (see Sections 1.2 and 1.3 for illustrations of this phenomenon).

For $a_0, \dots, a_{n-1} \in \alpha$, let $[a_0, \dots, a_{n-1}] = \bigcap_0^{n-1} T^{-i}(a_i)$. It is a *cylinder* of length n . For $f : X \rightarrow G$ and $Z \subset X$, set

$$Df(Z) = \inf\{C > 0 : \exists \Omega \subset Z \text{ with } m(Z \setminus \Omega) = 0 \text{ such that } \forall x, y \in \Omega, |f(x) - f(y)| \leq Cd(x, y)\}.$$

The main result of this note is the following theorem:

Theorem 1.1. *Let (X, T, m, α) be a probability preserving Gibbs-Markov map. Let $f : X \rightarrow G$ satisfy $\sum_{a \in \alpha} m(a)Df(a) < +\infty$. Let $u : X \rightarrow G$ be a measurable function such that $f = u - u \circ T$ almost everywhere.*

Then $\sup_{a_ \in \alpha_*} Du(a_*) < \infty$, where α_* is the partition generated by the images of the elements of α . Moreover, the function u is essentially bounded.*

Remarks:

- (1) Since T is Markov, α_* is coarser than α . In particular, $\sup_{a \in \alpha} Du(a) < \infty$, i.e., u has a version which is uniformly Lipschitz on each element of the partition α .
- (2) The map T is also Gibbs-Markov for the distance $d(x, y)^\gamma$ when $\gamma \in (0, 1]$. Hence, Theorem 1.1 implies a similar statement for Hölder functions.
- (3) The proof will in fact show that there exists a constant C depending only on T such that $\sup_{a_* \in \alpha_*} Du(a_*) \leq C \sum_{a \in \alpha} m(a)Df(a)$. In particular, when f is constant on each element of α , we get $Du(a_*) = 0$, i.e., u is essentially constant on the elements of α_* . When $G = S^1$, we get a completely different proof of [AD01, Theorem 3.1].

- (4) The proof would be easier under the stronger assumption

$$\sup_{a \in \alpha} Df(a) < \infty.$$

However, this assumption is too strong, since it is not compatible with the induction process which will enable us to extend Theorem 1.1 to non uniformly expanding settings.

In this paper, $\mathbb{N} = \{n \in \mathbb{Z}, n \geq 0\}$ and $\mathbb{N}^* = \mathbb{N} \setminus \{0\}$.

1.2. Application to Young towers. Let (X, d, m) be a probability space endowed with a bounded metric d . A map $T : X \rightarrow X$ is a *Young tower* ([You99]) if there exist integers $R_l \in \mathbb{N}^*$ and a partition $\{\Delta_{k,l}\}_{l \in \mathbb{N}, k \in \{0, \dots, R_l-1\}}$ of X such that

- (1) For all l and $k < R_l - 1$, T is a measurable isomorphism between $\Delta_{k,l}$ and $\Delta_{k+1,l}$, preserving m .
- (2) For all l , T is a measurable isomorphism between $\Delta_{R_l-1,l}$ and $\Delta_0 := \bigcup_m \Delta_{0,m}$.
- (3) There exists $\lambda > 1$ such that, for all l , for all $x, y \in \Delta_{0,l}$, $d(T^{R_l}x, T^{R_l}y) \geq \lambda d(x, y)$.
- (4) There exists $C > 0$ such that, for all l and $k < R_l$, for all $x, y \in \Delta_{k,l}$, $d(x, y) \leq Cd(T^{R_l-k}x, T^{R_l-k}y)$.
- (5) For $x \in \Delta_{R_l-1,l}$, let $g(x)$ be the inverse of the distortion of T at x , i.e., $g(x) = \frac{dm|_{\Delta_{R_l-1,l}}}{d(m \circ T|_{\Delta_{R_l-1,l}})}(x)$. There exists $C > 0$ such that, for all l , for all $x, y \in \Delta_{R_l-1,l}$, $\left|1 - \frac{g(x)}{g(y)}\right| \leq Cd(Tx, Ty)$.

The third and fifth conditions mean that the returns to the basis are expanding and have a controlled distortion. Hence, Young towers are a good model for many non uniformly expanding maps: the map has good properties, but after some waiting time which can be arbitrarily long.

Theorem 1.2. *Let (X, T, m, d) be a Young tower, and let $f : X \rightarrow G$ satisfy*

$$\sum m(\Delta_{k,l}) Df(\Delta_{k,l}) < \infty.$$

If $u : X \rightarrow G$ is such that $f = u - u \circ T$ almost everywhere, then the function u has a version which is Lipschitz on Δ_0 , i.e., there exists $C > 0$ such that, for almost all $x, y \in \Delta_0$, $|u(x) - u(y)| \leq Cd(x, y)$.

This result applies in particular when the function f is Lipschitz.

Proof. By [You99], we can assume without loss of generality that m is invariant.

Let $Y = \Delta_0$ with the partition $\alpha = \{\Delta_{0,l}\}$, $\varphi : Y \rightarrow \mathbb{N}^*$ the first return time to Y (i.e., on $\Delta_{0,l}$, $\varphi = R_l$), and $T_Y = T^\varphi$ the map induced by T on Y . Define also a distance d' on $\Delta_{0,l} \in \alpha$ by $d'(x, y) = d(T^{R_l}x, T^{R_l}y)$. If x and y are in two different elements of the partition α , set also $d'(x, y) = \lambda \sup_{X \times X} d$. Then $(Y, T_Y, m_Y/m(Y), d')$ is a Gibbs-Markov map for the partition α . Moreover, T_Y preserves the measure $m_Y/m(Y)$ and the partition α_* is the trivial partition.

Let $f : X \rightarrow G$ satisfy $\sum m(\Delta_{k,l})Df(\Delta_{k,l}) < \infty$, and assume that $f = u - u \circ T$. Define a function f_Y on Y by $f_Y(x) = \sum_{k=0}^{\varphi(x)-1} f(T^k x)$. On $\Delta_{0,l}$,

$$\begin{aligned} |f_Y(x) - f_Y(y)| &\leq \sum_{k=0}^{R_l-1} |f(T^k x) - f(T^k y)| \leq \sum_{k=0}^{R_l-1} d(T^k x, T^k y) Df(\Delta_{k,l}) \\ &\leq Cd(T^{R_l} x, T^{R_l} y) \sum_{k=0}^{R_l-1} Df(\Delta_{k,l}) = Cd'(x, y) \sum_{k=0}^{R_l-1} Df(\Delta_{k,l}). \end{aligned}$$

Hence, $\sum_{a \in \alpha} m(a)Df_Y(a) \leq C \sum m(\Delta_{k,l})Df(\Delta_{k,l}) < \infty$. Moreover, $f_Y = u - u \circ T_Y$.

Theorem 1.1 applies and proves that u is almost everywhere Lipschitz on each element of α_* , for the distance d' . In particular, on any element $\Delta_{0,l}$ of α , we get $|u(x) - u(y)| \leq Ed'(x, y)$.

Take finally $x', y' \in \Delta_0$. They have preimages x, y under T^{R_l} in $\Delta_{0,l}$. As $f_Y(x) = u(x) - u(x')$ and $f_Y(y) = u(y) - u(y')$, we get

$$\begin{aligned} |u(x') - u(y')| &\leq |f_Y(x) - f_Y(y)| + |u(x) - u(y)| \leq Cd'(x, y) + Ed'(x, y) \\ &= C'd(x', y'). \end{aligned} \quad \square$$

1.3. Applications to intermittent maps. For $\alpha \in (0, 1)$, let T be the map from $[0, 1]$ to itself given by

$$T(x) = \begin{cases} x(1 + 2^\alpha x^\alpha) & \text{if } 0 \leq x \leq 1/2; \\ 2x - 1 & \text{if } 1/2 < x \leq 1. \end{cases}$$

This map has been studied by [LSV99]. It is nonuniformly expanding since the fixed point 0 satisfies $T'(0) = 1$, and admits an absolutely continuous invariant probability measure.

Proposition 1.3. *Let $f : [0, 1] \rightarrow G$ be Hölder with exponent $\gamma > 0$ on the intervals $[0, 1/2]$ and $(1/2, 1]$. If $u : [0, 1] \rightarrow G$ is measurable and satisfies $f = u - u \circ T$ Lebesgue almost everywhere, then there exists a function $\tilde{u}$, equal to u almost everywhere, Hölder with exponent γ , and such that $f = \tilde{u} - \tilde{u} \circ T$ everywhere.*

Proof. Let $Y = (1/2, 1]$, φ the first return time from Y to itself and $T_Y : Y \rightarrow Y$ the induced map. Then T_Y is Gibbs-Markov for the partition $B_n = \{y \in Y : \varphi(y) = n\}$, by [LSV99]. Hence, the arguments in the proof of Theorem 1.2 apply and prove that u is a.e. Hölder on Y . As $T : (1/2, 1] \rightarrow (0, 1]$ is Lipschitz and has Lipschitz inverse, the coboundary equation implies that u is a.e. Hölder on $(0, 1]$, i.e., there exists a set V of full measure and a constant C such that, for all $x, y \in V$, $|u(x) - u(y)| \leq C|x - y|^\gamma$.

The function u is uniformly continuous on V , whence it can be extended to a continuous – and even Hölder – function $\tilde{u}$ on $[0, 1]$. On $V \cap T^{-1}(V)$, which is dense, we have $f(x) = \tilde{u}(x) - \tilde{u}(Tx)$. Since both members of this equality are continuous on the intervals $[0, 1/2]$ et $(1/2, 1]$, this equality holds in fact everywhere. $\square$

In particular, if f is a measurable coboundary, it satisfies $\sum_{k=0}^{n-1} f(T^k x) = \tilde{u}(T^n x) - \tilde{u}(x) = 0$ at any point x such that $T^n(x) = x$.

Corollary 1.4. *If $f : [0, 1] \rightarrow \mathbb{R}$ is Hölder continuous on $[0, 1/2]$ and $(1/2, 1]$ and satisfies $f(0) \neq 0$, then f is not a measurable coboundary.*

This solves a conjecture stated in [FHV03]: in this article, the authors need to know that $f = \log |T'| - \int \log |T'|$ is not a coboundary to get a nonzero variance in the central limit theorem. As f is α -Hölder on $[0, 1/2]$ and $(1/2, 1]$, and $f(0) = -\int \log |T'| < 0$, the corollary applies and proves that it is indeed never the case.

Using Theorem 1.1 with $G = S^1$, we can get in the same way a stronger result:

Corollary 1.5. *The function $f(x) = \log |T'| - \int \log |T'|$ can not be written as $f = u - u \circ T + \lambda q + \mu$ almost everywhere, where $u : [0, 1] \rightarrow \mathbb{R}$ is measurable, $q : X \rightarrow \mathbb{Z}$ and $\lambda, \mu \in \mathbb{R}$.*

The proof is the same, using the behavior at the fixed points 0 and 1 to get a contradiction. This is a strong aperiodicity result on the function f . By [Gou03, Theorem 1.2], it implies that f satisfies a local limit theorem when $\alpha < 1/2$.

1.4. Application to positive recurrent Markov shifts. Let $T : X \rightarrow X$ be a positive recurrent Markov shift with Hölder potential, as defined in [Sar01], preserving the probability measure m . The map T satisfies the same assumptions as a Gibbs-Markov map, except the BIP property. We assume also that the distance d is given by $d(x, y) = \tau^{s(x, y)}$ where $\tau \in (0, 1)$ and $s(x, y)$ is the separation time of x and y . Such maps have in general more complicated combinatorics than Young towers, but they enjoy uniform expansion (since $d(Tx, Ty) = \tau^{-1}d(x, y)$ for all x, y in the same element of α) while Young towers are expanding only after many iterates.

Theorem 1.6. *Let $f : X \rightarrow G$ satisfy $\sum_{a \in \alpha} m(a)Df(a) < \infty$. Let $u : X \rightarrow G$ be a measurable function such that $f = u - u \circ T$ almost everywhere. Then, for all $a \in \alpha$, $Du(a) < \infty$. Moreover, if T is transitive, $\sum_{a \in \alpha} m(a)Du(a) < \infty$.*

Proof. For $a \in \alpha$, let T_a be the map induced by T on $[a]$. It is Gibbs-Markov. Using Theorem 1.1, we show as in the proof of Theorem 1.2 that $Du(a) < \infty$. If T is transitive, the proof of Lemma 2.3 applies and gives $\sum m(a)Du(a) < \infty$. $\square$

2. PROOF OF THEOREM 1.1

A Gibbs-Markov map is *transitive* if, for all $a, b \in \alpha$, there exists n such that $b \subset T^n(a) \bmod 0$. When T preserves a probability measure, there exists a finite decomposition $\alpha = \alpha_1 \cup \dots \cup \alpha_n$ such that the image of an element of α_i is contained in $X_i = \bigcup_{a \in \alpha_i} a$, and such that T is a transitive Gibbs-Markov map on X_i ([Aar97]). To prove the theorem, it is sufficient to prove it on each X_i . We can therefore assume that T is transitive.

The main step of the proof is the following lemma:

Lemma 2.1. *There exists $\alpha_1 \in \alpha$ such that $Du(\alpha_1) < \infty$.*

Proof. Let $\Phi(x) = Df(a)$ when $x \in a$. This function is integrable by assumption. In particular, there exists a set X_1 of full measure such that the Birkhoff sums $S_n \Phi(x) = \sum_{k=0}^{n-1} \Phi(T^k x)$ satisfy $S_n \Phi(x) = O(n)$ when $x \in X_1$.

There exists X_2 of full measure such that, if $x \in X_2$, all its iterates satisfy: for almost all y in the same element of partition a as $T^n x$, $|f(y) - f(T^n x)| \leq Df(a)d(y, T^n x)$.

The martingale convergence theorem implies that almost every point is a measurable continuity point of u : there exists X_3 of full measure such that, if $x \in X_3$ and

$a_0, a_1, \dots$ denotes the sequence of elements of α containing respectively $x, Tx, \dots$, then, for all $\varepsilon > 0$,

$$\frac{m\{y \in [a_0, \dots, a_{n-1}] : |u(y) - u(x)| > \varepsilon\}}{m[a_0, \dots, a_{n-1}]} \rightarrow 0.$$

As T is Gibbs-Markov, all its iterates have a bounded distortion ([Aar97, Proposition 4.3.1]). Hence, there exists $B > 0$ such that, for any measurable set Z and for any cylinder of length k ,

$$(2) \quad B^{-1} \frac{m(T(a_{k-1}) \cap Z)}{m(Ta_{k-1})} \leq \frac{m([a_0, \dots, a_{k-1}] \cap T^{-k}Z)}{m[a_0, \dots, a_{k-1}]} \leq B \frac{m(T(a_{k-1}) \cap Z)}{m(Ta_{k-1})}.$$

Since T has the big image property, this implies that there exists $B' > 0$ such that

$$(3) \quad \frac{m([a_0, \dots, a_{k-1}] \cap T^{-k}Z)}{m[a_0, \dots, a_{k-1}]} \leq B' m(Z).$$

Let $\lambda > 1$ be the expansion factor of T and let $K > 0$ be large enough so that

$$(4) \quad K \log \lambda > 3.$$

Let $\alpha_1, \dots, \alpha_N$ be a finite number of elements of α such that $m(X \setminus \bigcup \alpha_i) \leq \varepsilon_0$ where ε_0 satisfies $K \log(1 - B'\varepsilon_0) \geq -1/2$. Write

$$Z_n = \{x : \forall n^3 \leq k < n^3 + \lfloor K \log n \rfloor, T^k(x) \in \alpha_1 \cup \dots \cup \alpha_N\}.$$

Let finally X_4 be the set of points belonging to infinitely many Z_n .

Lemma 2.2. *The set X_4 has nonzero measure.*

Proof. Write $A = \alpha_1 \cup \dots \cup \alpha_N$. Let us first bound $m(Z_n)$ from below. For any cylinder $[a_0, \dots, a_{k-1}]$, we apply (3) to $X \setminus A$, of measure at most ε_0 , and we get

$$m([a_0, \dots, a_{k-1}] \cap T^{-k}A) \geq (1 - B'\varepsilon_0)m[a_0, \dots, a_{k-1}].$$

Summing these inequalities for $a_{k-1} = \alpha_1, \dots, \alpha_N$ yields

$$m([a_0, \dots, a_{k-2}] \cap T^{-k+1}A \cap T^{-k}A) \geq (1 - B'\varepsilon_0)m([a_0, \dots, a_{k-2}] \cap T^{-k+1}A).$$

This last term is larger than $(1 - B'\varepsilon_0)^2 m[a_0, \dots, a_{k-2}]$, again by (3). We get in this way by induction

$$m([a_0, \dots, a_l] \cap T^{-l-1}A \cap \dots \cap T^{-k}A) \geq (1 - B'\varepsilon_0)^{k-l} m[a_0, \dots, a_l].$$

In particular, for $l = -1$ and $k = \lfloor K \log n \rfloor - 1$, we get using the invariance of m that

$$m(Z_n) \geq (1 - B'\varepsilon_0)^{K \log n} = n^{K \log(1 - B'\varepsilon_0)} \geq \frac{1}{\sqrt{n}}.$$

Hence, $\sum m(Z_n) = \infty$. We will use a version of the Borel-Cantelli Lemma to conclude. Since the sets Z_n are not independent, we will use the following version of this lemma, due to Lamperti ([Spi64, Proposition 6.26.3]):

If $\sum m(Z_n) = \infty$ and

$$\liminf_{n \rightarrow \infty} \frac{\sum_{j,k=1}^n m(Z_j \cap Z_k)}{(\sum_{k=1}^n m(Z_k))^2} < \infty$$

then the set of points belonging to infinitely many Z_n has nonzero measure.

To estimate $m(Z_j \cap Z_k)$, we will use the transfer operator $\hat{T}$, defined on L^2 as the adjoint of the composition by T . It acts continuously on the space $\mathcal{L}$ of

functions which are bounded and Lipschitz on any element of α . Moreover, by [Aar97, Proposition 4.7.3], there exist $M > 0$ and $\eta < 1$ such that, for any $h \in \mathcal{L}$,

$$(5) \quad \|\widehat{T}^p h\|_{\mathcal{L}} \leq M(\eta^p \|h\|_{\mathcal{L}} + \|h\|_1).$$

Let χ be the characteristic function of A , and $\gamma_n = \prod_{0 \leq k < \lfloor K \log n \rfloor} \chi \circ T^k$: hence, $m(Z_n) = \int \gamma_n \circ T^{n^3} = \int \gamma_n$, and $m(Z_n \cap Z_p) = \int \gamma_n \circ T^{n^3} \cdot \gamma_p \circ T^{p^3}$. The function χ belongs to $\mathcal{L}$. For $k > j$,

$$(6) \quad m(Z_j \cap Z_k) = \int \gamma_j \circ T^{j^3} \cdot \gamma_k \circ T^{k^3} = \int \widehat{T}^{k^3-j^3}(\gamma_j) \cdot \gamma_k \leq \|\widehat{T}^{k^3-j^3}(\gamma_j)\|_{\mathcal{L}} m(Z_k).$$

As $\widehat{T}$ acts continuously on $\mathcal{L}$, the function

$$\delta_j = \widehat{T}^{\lfloor K \log j \rfloor}(\gamma_j) = \widehat{T}(\chi \widehat{T}(\chi \cdots \widehat{T}(\chi) \cdots))$$

satisfies $\|\delta_j\|_{\mathcal{L}} \leq (2M)^{K \log j}$. The inequality (5) applied to $p = k^3 - j^3 - \lfloor K \log j \rfloor$ and $h = \delta_j$ yields

$$(7) \quad \begin{aligned} \|\widehat{T}^{k^3-j^3} \gamma_j\|_{\mathcal{L}} &\leq M \left(\eta^{k^3-j^3-K \log j} \|\delta_j\|_{\mathcal{L}} + \|\delta_j\|_1 \right) \\ &\leq M \left(\eta^{k^3-j^3-K \log j} (2M)^{K \log j} + m(Z_j) \right) \end{aligned}$$

since $\|\delta_j\|_1 = \int \delta_j = \int \gamma_j$, for all these functions are nonnegative. Hence, (6) and (7) give

$$|m(Z_j \cap Z_k)| \leq M \eta^{k^3-j^3} (2M/\eta)^{K \log j} + M m(Z_j) m(Z_k).$$

Finally,

$$\begin{aligned} \sum_{j < k \leq n} m(Z_j \cap Z_k) &\leq M \sum_{j < k} m(Z_j) m(Z_k) + M \sum_{j=1}^{\infty} \eta^{-j^3} (2M/\eta)^{K \log j} \sum_{k=j+1}^{\infty} \eta^{k^3} \\ &\leq M \left(\sum_{k \leq n} m(Z_k) \right)^2 + M \sum_{j=1}^{\infty} \eta^{-j^3} (2M/\eta)^{K \log j} \sum_{l=(j+1)^3}^{\infty} \eta^l. \end{aligned}$$

The last sum is bounded by

$$M \sum_{j=1}^{\infty} \eta^{-j^3} (2M/\eta)^{K \log j} \frac{\eta^{(j+1)^3}}{1-\eta} < \infty,$$

which shows that the aforementioned Borel-Cantelli lemma applies. $\square$

We can take $x_0 \in X_1 \cap X_2 \cap X_3 \cap X_4$ since this set has positive measure. Let $m_k \rightarrow \infty$ be such that $x_0 \in Z_{m_k}$, and $n_k = m_k^3 + \lfloor K \log m_k \rfloor - 1$. Then $T^{n_k}(x_0)$ belongs to one of the sets $\alpha_1, \dots, \alpha_N$. In particular, one of these sets is used infinitely many times, and taking a further subsequence we can for example assume that $T^{n_k}(x_0) \in \alpha_1$ for all k . We will show that $Du(\alpha_1) < \infty$. Denote by $a_0, a_1, \dots$ the elements of α containing respectively $x_0, T(x_0), \dots$. Let $[\underline{a}_n] = [a_0, \dots, a_{n-1}]$, and let $v_n : Ta_{n-1} \rightarrow [\underline{a}_n]$ be the inverse of $T^n : [\underline{a}_n] \rightarrow Ta_{n-1}$.

Let $\varepsilon > 0$. As $x_0 \in X_3$,

$$\frac{m\{y \in [\underline{a}_{n_k}] : |u(y) - u(x_0)| > \varepsilon\}}{m[\underline{a}_{n_k}]} \rightarrow 0.$$

Taking a further subsequence of n_k , we can assume that

$$\sum \frac{m\{y \in [\underline{a}_{n_k}] : |u(y) - u(x_0)| > \varepsilon\}}{m[\underline{a}_{n_k}]} < \infty.$$

For all $k \in \mathbb{N}$, the distortion control (2) implies that

$$\frac{m\{y \in Ta_{n_k-1} : |u(v_{n_k}y) - u(x_0)| > \varepsilon\}}{m[Ta_{n_k-1}]} \prec \frac{m\{y \in [\underline{a}_{n_k}] : |u(y) - u(x_0)| > \varepsilon\}}{m[\underline{a}_{n_k}]}.$$

Hence, $\sum_k m\{y \in Ta_{n_k-1} : |u(v_{n_k}y) - u(x_0)| > \varepsilon\} < +\infty$. Therefore, $U_\varepsilon := \{y \in X : \exists \kappa, \forall k \geq \kappa, \text{ if } y \in Ta_{n_k-1} \text{ then } |u(v_{n_k}y) - u(x_0)| \leq \varepsilon\}$ has full measure.

Let $y_1, y_2 \in U_\varepsilon \cap \alpha_1$. If k is large enough, the preimages y'_1 and y'_2 of y_1 and y_2 under T^{n_k} in $[\underline{a}_{n_k}]$ satisfy $|u(y'_i) - u(x_0)| \leq \varepsilon$, whence $|u(y'_1) - u(y'_2)| \leq 2\varepsilon$. Then

$$\begin{aligned} |u(y_1) - u(y_2)| &= |u \circ T^{n_k}(y'_1) - u \circ T^{n_k}(y'_2)| \\ (8) \quad &\leq \sum_{i=0}^{n_k-1} |f \circ T^i(y'_1) - f \circ T^i(y'_2)| + |u(y'_1) - u(y'_2)|. \end{aligned}$$

Recall that $n_k = m_k^3 + \lfloor K \log m_k \rfloor - 1$, and that Φ is defined by $\Phi(x) = Df(a)$ when $x \in a$. Then

$$\begin{aligned} (9) \quad \sum_{i=0}^{m_k^3-1} |f \circ T^i(y'_1) - f \circ T^i(y'_2)| &\leq \sum_{i=0}^{m_k^3-1} \Phi(T^i(x_0)) d(T^i y'_1, T^i y'_2) \\ &\leq \sum_{i=0}^{m_k^3-1} \Phi(T^i(x_0)) \lambda^{i-n_k} d(T^{n_k} y'_1, T^{n_k} y'_2) \\ &\leq \lambda^{-K \log m_k + 2} S_{m_k^3} \Phi(x_0) d(y_1, y_2). \end{aligned}$$

Since $x_0 \in X_1$, there exists C such that $S_n \Phi(x_0) \leq Cn$ for all n . As $-K \log \lambda < -3$ by (4), we get that (9) tends to 0.

Finally, set $D = \sup Df(\alpha_j)$ for $1 \leq j \leq N$. By definition of m_k , we have $T^i(x_0) \in \bigcup_{1 \leq j \leq N} \alpha_j$ for all $m_k^3 \leq i < n_k$, whence

$$\begin{aligned} \sum_{i=m_k^3}^{n_k-1} |f \circ T^i(y'_1) - f \circ T^i(y'_2)| &\leq \sum_{i=m_k^3}^{n_k-1} D d(T^i y'_1, T^i y'_2) \leq D \sum_{i=m_k^3}^{n_k-1} \lambda^{i-n_k} d(y_1, y_2) \\ &\leq \frac{D}{\lambda-1} d(y_1, y_2). \end{aligned}$$

Equation (8) then yields

$$|u(y_1) - u(y_2)| \leq o(1) + \frac{D}{\lambda-1} d(y_1, y_2) + 2\varepsilon.$$

Finally, on $\alpha_1 \cap \bigcap_{\varepsilon > 0} U_\varepsilon$, we have $|u(y_1) - u(y_2)| \leq \frac{D}{\lambda-1} d(y_1, y_2)$. $\square$

Lemma 2.3. *We have $\sum_{a \in \alpha} m(a) Du(a) < \infty$.*

Proof. Let us show that, for any $a \in \alpha$, $Du(a) < \infty$. As T is transitive, there exists n such that $a \subset T^n(\alpha_1)$. Let $[a_0, \dots, a_{n-1}]$ be a cylinder included in α_1 such

that $a \subset T(a_{n-1})$. For $y_1, y_2 \in a$, let y'_1 and y'_2 be their preimages under T^n in $[a_0, \dots, a_{n-1}]$. Then

$$(10) \quad \begin{aligned} |u(y_1) - u(y_2)| &\leq \sum_{i=0}^{n-1} |f(T^i y'_1) - f(T^i y'_2)| + |u(y'_1) - u(y'_2)| \\ &\leq \sum_{i=0}^{n-1} Df(a_i) \lambda^{i-n} d(y_1, y_2) + \lambda^{-n} Du(\alpha_1) d(y_1, y_2), \end{aligned}$$

which proves that $Du(a) < \infty$.

Let β be a finite nonempty subset of α . For $a \in \alpha \setminus \beta$, let us show

$$(11) \quad m(a) = \sum_{n=1}^{\infty} \sum_{a_0 \in \beta, a_1, \dots, a_{n-1} \in \alpha \setminus \beta} m[a_0, a_1, \dots, a_{n-1}, a].$$

Let $Y = \bigcup_{b \in \beta} b$. Write $A_0 = a$, and $A_{n+1} = T^{-1}(A_n) \setminus Y$ and $B_{n+1} = T^{-1}(A_n) \cap Y$. We get

$$A_n = \bigcup_{a_0, \dots, a_{n-1} \in \alpha \setminus \beta} [a_0, \dots, a_{n-1}, a] \text{ and } B_n = \bigcup_{a_0 \in \beta, a_1, \dots, a_{n-1} \in \alpha \setminus \beta} [a_0, \dots, a_{n-1}, a].$$

Thus, we want to show that $m(a) = \sum_n m(B_n)$. The equality $T^{-1}(A_n) = A_{n+1} \cup B_{n+1}$ implies $m(A_n) = m(A_{n+1}) + m(B_{n+1})$. By induction, we get $m(a) = m(B_1) + \dots + m(B_n) + m(A_n)$. It remains to prove that $m(A_n) \rightarrow 0$. Note that $A_n \subset C_n = \{x : \forall 0 \leq k \leq n, T^k(x) \notin Y\}$. We will show that $m(C_n) \rightarrow 0$ by proving that $C = \bigcap C_n$ has 0 measure. Since the measure is invariant and $C \subset T^{-1}(C)$, $C = T^{-1}(C) \pmod{0}$, whence $m(C) = 0$ or 1 by ergodicity ([Aar97, Theorem 4.4.7]). The set C does not intersect Y , which has nonzero measure, hence $m(C) = 0$. This proves (11).

Let $[a_0, \dots, a_{n-1}, a]$ be a cylinder of nonzero measure. By (10),

$$Du(a) \leq \sum_{i=0}^{n-1} \lambda^{i-n} Df(a_i) + \lambda^{-n} Du(a_0).$$

Hence, (11) yields

$$\begin{aligned} m(a) Du(a) &\leq \sum_{n=1}^{\infty} \sum_{a_0 \in \beta, a_1, \dots, a_{n-1} \in \alpha \setminus \beta} m[a_0, \dots, a_{n-1}, a] \left(\sum_{i=0}^{n-1} \lambda^{i-n} Df(a_i) \right) \\ &\quad + m(a) \sup_{b \in \beta} Du(b). \end{aligned}$$

As $\sum m(a) \sup_{b \in \beta} Du(b) < \infty$, we will show that $\sum m(a) Du(a) < \infty$ by showing that

$$(12) \quad \sum_{n=1}^{\infty} \sum_{a_0 \in \beta, a_1, \dots, a_{n-1} \in \alpha \setminus \beta} m[a_0, \dots, a_{n-1}] \left(\sum_{i=0}^{n-1} \lambda^{i-n} Df(a_i) \right)$$

is finite. In this expression, for $a' \in \alpha \setminus \beta$, the prefactor of a term $\lambda^{-k} Df(a')$ is

$$\begin{aligned} \sum_{n=1}^{\infty} \sum_{\substack{a_0 \in \beta, a_1, \dots, a_{n-1} \in \alpha \setminus \beta \\ a_{n+1}, \dots, a_{n+k-1} \in \alpha \setminus \beta}} m[a_0, \dots, a_{n-1}, a', a_{n+1}, \dots, a_{n+k-1}] \\ \leq \sum_{n=1}^{\infty} \sum_{a_0 \in \beta, a_1, \dots, a_{n-1} \in \alpha \setminus \beta} m[a_0, \dots, a_{n-1}, a']. \end{aligned}$$

By (11), this last term is equal to $m(a')$. In (12), the prefactor of a term $\lambda^{-k} Df(a')$ with $a' \in \beta$ is also at most $m(a')$. Hence,

$$(12) \leq \sum_{a' \in \alpha} \sum_{k=1}^{\infty} m(a') \lambda^{-k} Df(a'),$$

which is finite since $\sum m(a') Df(a') < \infty$. $\square$

Proof of Theorem 1.1. For almost all x , $\sum_{T y=x} g(y) = 1$. Let us write $T^{-1}(x) = \{x_0, x_1, \dots\}$, and let a_i be the element of α containing x_i . By bounded distortion and the big image property, there exists $C > 0$ such that, for all n , $g(x_n) \leq C m(a_n)$. As $\sum g(x_n) = 1$, this implies $C \sum m(a_n) \geq 1$.

Let a_* be an element of α_* . Let $x, y \in a_*$. By definition of α_* , their preimages $x_0, x_1, \dots$ and $y_0, y_1, \dots$ belong to the same elements $a_0, a_1, \dots$ of α . Since $f = u - u \circ T$, we have for any n

$$\begin{aligned} |u(x) - u(y)| &\leq |f(x_n) - f(y_n)| + |u(x_n) - u(y_n)| \leq (Df(a_n) + Du(a_n)) d(x_n, y_n) \\ &\leq (Df(a_n) + Du(a_n)) \lambda^{-1} d(x, y). \end{aligned}$$

Hence,

$$\begin{aligned} |u(x) - u(y)| &\leq C \sum m(a_n) |u(x) - u(y)| \\ &\leq C \sum m(a_n) (Df(a_n) + Du(a_n)) \lambda^{-1} d(x, y). \end{aligned}$$

Finally, $Du(a_*) \leq \frac{C}{\lambda} \sum_{a \in \alpha} m(a) (Df(a) + Du(a))$, which is finite by Lemma 2.3.

To prove that u is essentially bounded, we use the big preimage property. Let $a_1, \dots, a_n \in \alpha$ be such that every element of α is contained in the image of some a_i . Let $a \in \alpha$, and let i be such that $a \subset T(a_i)$. For $x \in a$, let x' be its preimage in a_i , we get

$$|u(x)| = |u(x') - f(x')| \leq \|u|_{a_i}\|_{\infty} + \|f|_{a_i}\|_{\infty}.$$

This last quantity is uniformly bounded. $\square$

REFERENCES

- [Aar97] Jon Aaronson. *An introduction to infinite ergodic theory*, volume 50 of *Mathematical Surveys and Monographs*. American Mathematical Society, 1997.
- [AD01] Jon Aaronson and Manfred Denker. Local limit theorems for partial sums of stationary sequences generated by Gibbs-Markov maps. *Stoch. Dyn.*, 1:193–237, 2001.
- [ADSZ04] Jon Aaronson, Manfred Denker, Omri Sarig, and Roland Zweimüller. Aperiodicity of cocycles and conditional local limit theorems. *Stoch. Dyn.*, 4:31–62, 2004.
- [FHV03] Pierre Ferrero, Nicolai T.A. Haydn, and Sandro Vaienti. Entropy fluctuations for parabolic maps. *Nonlinearity*, 16:1203–1218, 2003.
- [GH88] Yves Guivarc'h and Jean Hardy. Thormes limites pour une classe de chaines de Markov et applications aux difféomorphismes d'Anosov. *Ann. Inst. H. Poincaré Probab. Statist.*, 24:73–98, 1988.

- [Gou03] Sébastien Gouëzel. Berry-Esseen theorem and local limit theorem for non uniformly expanding maps. Preprint, 2003.
- [Leo60] V. P. Leonov. On the central limit theorem for ergodic endomorphisms of compact commutative groups. *Dokl. Akad. Nauk SSSR*, 135:258–261, 1960.
- [Liv72] Alexander N. Livšic. Cohomology properties of dynamical systems. *Math. USSR Izv.*, pages 1278–1301, 1972.
- [Liv96] Carlangelo Liverani. Central limit theorems for deterministic systems. In *International Conference on Dynamical Systems, Montevideo 1995*, volume 362 of *Pitman Research Notes in Mathematics*, 1996.
- [LSV99] Carlangelo Liverani, Benoît Saussol, and Sandro Vaienti. A probabilistic approach to intermittency. *Ergodic Theory Dynam. Systems*, 19:671–685, 1999.
- [NS03] Matthew Nicol and Andrew Scott. Livšic theorems and stable ergodicity for group extensions of hyperbolic systems with discontinuities. *Ergodic Theory Dynam. Systems*, 23:1867–1889, 2003.
- [PY99] Mark Pollicott and Michiko Yuri. Regularity of solutions to the measurable Livsic equation. *Trans. Amer. Math. Soc.*, 351:559–568, 1999.
- [Sar01] Omri Sarig. Thermodynamic formalism for null recurrent potentials. *Israel J. Math.*, 121:285–311, 2001.
- [Sar03] Omri Sarig. Existence of Gibbs measures for countable Markov shifts. *Proc. Amer. Math. Soc.*, 131(6):1751–1758, 2003.
- [Spi64] Frank Spitzer. *Principles of random walk*. The University Series in Higher Mathematics. D. Van Nostrand Co., Inc., Princeton, N.J.-Toronto-London, 1964.
- [You99] Lai-Sang Young. Recurrence times and rates of mixing. *Israel J. Math.*, 110:153–188, 1999.

DÉPARTEMENT DE MATHÉMATIQUES ET APPLICATIONS, ÉCOLE NORMALE SUPÉRIEURE, 45 RUE D'ULM, 75005 PARIS, FRANCE.

E-mail address: `sebastien.gouezel@ens.fr`